

# Head and Neck Cancer May be Sexually Transmitted; CBCD Reviews a Study

*There is an association between HPV, sexual behavior, and head and neck cancer. (1)*

ROCHESTER, NEW YORK, UNITED STATES OF AMERICA, September 24, 2015 /EINPresswire.com/ --

“Infected with the human papillomavirus (HPV)? The CBCD recommends taking [Gene-Eden-VIR](#) or [Novirin](#).” - Greg Bennett, CBCD

Head and neck cancers are associated with oral sexual behavior, according to multiple studies. For example, Dr. Eleni Rettig and colleagues wrote that “HPV-positive oropharyngeal squamous cell carcinoma (HPV-OSCC) is associated with oral sexual behaviors.” (2) Dr. Rettig is from the Department of Otolaryngology-Head and Neck Surgery at Johns Hopkins Medicine. Research shows that worldwide rates of oral HPV infection and resulting head and neck cancers are rising. (3) Many researchers believe that this is the result of changing attitudes about oral sex. In the past, this practice was considered slightly taboo, but this belief has changed over the past 50 years. As Dr. Rettig wrote, “the sharp rise in incidence of HPV-OSCC in the USA has been attributed to changes in sexual norms over the past five decades, with lower age at sexual debut and higher numbers of sexual partners per individual.” (2)



The Center for the Biology of Chronic Disease (CBCD) points out that, in addition to cancer, oral HPV can also cause warts in the mouth. Dr. Percinoto and colleagues wrote that genital warts in the mouth are a “sexually transmitted disease (STD) appearing most frequently as soft, pink cauliflower like growths in moist areas, such as the genitalia, mouth and other places. The disease is highly contagious, (and) can appear singly or in groups, small or large.” (4) Dr. Percinoto and colleagues are from the Aracatuba Dental School at the University of Estadual

infection is usually transmitted by sexual intercourse.” (4) The CBCD therefore recommends that anyone who suspects that they have an oral HPV infection talk to their doctors about Gene-Eden-VIR or Novirin. The formula of these natural antiviral products was shown to safely and effectively reduce symptoms associated with HPV in two separate post-marketing clinical studies that followed FDA guidelines.

Click to learn more about [HPV symptoms](#).

The formula of Gene-Eden-VIR and Novirin was tested by Hanan Polansky and Edan Itzkovitz from the CBCD in two clinical studies that followed FDA guidelines. The studies showed that the Novirin and Gene-Eden-VIR formula is effective against the HPV and other viruses. The clinical studies were published in the peer reviewed, medical journal Pharmacology & Pharmacy, the first, in a special edition on Advances in Antiviral Drugs. Study authors wrote that, “individuals infected with the HPV...reported a safe decrease in their symptoms following treatment with Gene-Eden-VIR.” (5) The study authors also wrote that “we observed a statistically significant decrease in the severity, duration, and frequency of symptoms.” (5)

Both products can be ordered online on the Gene-Eden-VIR and Novirin websites, here:

<http://www.gene-eden-vir.com>

and

<http://www.novirin.com>

Novirin and Gene-Eden-VIR are natural antiviral dietary supplements. Their formula contains five natural ingredients: Selenium, Camellia Sinesis Extract, Quercetin, Cinnamomum Extract, and Licorice Extract. The first ingredient is a trace element, and the other four are plant extracts. Each ingredient and its dose was chosen through a scientific approach. Scientists at polyDNA, the company that invented and patented the formula, scanned thousands of scientific and medical papers published in various medical and scientific journals, and identified the safest and most effective natural ingredients against latent viruses. To date, Novirin and Gene-Eden-VIR are the only natural antiviral products on the market with published clinical studies that support their claims.

The CDC notes that “some studies suggest that oral HPV may be passed on during oral sex (from mouth-to-genital or mouth-to-anus contact) or open-mouthed (“French”) kissing, others have not. The likelihood of getting HPV from kissing or having oral sex with someone who has HPV is not known. We do know that partners who have been together a long time tend to share genital HPV-meaning they both may have it.” (6) Additionally, the CDC says that “there is no FDA-approved test to diagnose HPV in the mouth or throat. Medical and dental organizations do not recommend screening for oral HPV.” Thus, there is no way to know for sure if a person has an oral HPV infection unless the person develops genital warts in the mouth.

Are there treatments available against the HPV?

“There are no drugs approved against the HPV. Current treatments include procedures, such as cryotherapy, conization, and the Loop Electrosurgical Excision Procedure (LEEP). These procedures use liquid nitrogen, a surgical knife (scalpel), a carbon dioxide (CO2) laser, or electrical current to remove the abnormal growths caused by the HPV. These growths include cells that harbor the active virus. The procedures do not target cells with the latent virus. Since they do not remove the latent virus, these procedures only produce a temporary remission.” (5) In contrast, Novirin and Gene-Eden-VIR were designed to target the latent HPV.

Find us on Facebook here: <https://www.facebook.com/GeneEdenVIR>

Follow us Twitter here: @HananPolansky

All orders of Gene-Eden-VIR and Novirin are completely confidential, and no information is shared or sold to any third party. Privacy is assured.

References:

(1) Farsi NJ1, El-Zein M2, Gaied H3, Lee YC4, Hashibe M5, Nicolau B3, Rousseau MC6. "Sexual behaviours and head and neck cancer: A systematic review and meta-analysis." *Cancer Epidemiol.* 2015 Sep 11. pii: S1877-7821(15)00174-5

(2) Eleni Rettig,1 Ana Ponce Kiess,2 and Carole Fakhry1,3,\* "The role of sexual behavior in head and neck cancer: implications for prevention and therapy" *Expert Rev Anticancer Ther.* Author manuscript; available in PMC 2015 Apr 5.

(3) McCarthy CE1, Field JK2, Rajlawat BP1, Field AE1, Marcus MW2. "Trends and regional variation in the incidence of head and neck cancers in England: 2002 to 2011." *Int J Oncol.* 2015 May 7. doi: 10.3892/ijo.2015.2990.

<http://www.ncbi.nlm.nih.gov/pubmed/25955390>

(4) Percinoto AC, Danelon M, Crivelini MM, Cunha RF, Percinoto C1. Condyloma acuminata in the tongue and palate of a sexually abused child: a case report. *BMC Res Notes.* 2014 Jul 23;7:467. [ncbi.nlm.nih.gov/pubmed/25053204](http://www.ncbi.nlm.nih.gov/pubmed/25053204)

(5) Polansky H, Itzkovitz E. Gene-Eden-VIR Is Antiviral: Results of a Post Marketing Clinical Study. *Pharmacology & Pharmacy*, 2013, 4, 1-8 <http://dx.doi.org/10.4236/pp.2013.46A001>

(6) CDC.org - Human papillomavirus (HPV) and Oropharyngeal Cancer - Fact Sheet. Last Updated on November 22, 2013.

[cdc.gov/std/hpv/stdfact-hpvandoropharyngealcancer.htm](http://cdc.gov/std/hpv/stdfact-hpvandoropharyngealcancer.htm)

Greg Bennett  
CBCD  
585-250-9999  
email us here

---

This press release can be viewed online at: <http://www.einpresswire.com>

Disclaimer: If you have any questions regarding information in this press release please contact the company listed in the press release. Please do not contact EIN Presswire. We will be unable to assist you with your inquiry. EIN Presswire disclaims any content contained in these releases.

© 1995-2015 IPD Group, Inc. All Right Reserved.





