

Warning: Acyclovir May Poison the Brain in Some Cases; The CBCD Reviews a Report

“Dosing recommended by the manufacturer's data sheet does not consider prevention of acyclovir-induced neurotoxicity.” (1)

ROCHESTER, NY, UNITED STATES,
December 3, 2015 /EINPresswire.com/ --

“Infected with the herpes virus (HSV-1 or HSV-2)? The CBCD recommends taking [Gene-Eden-VIR](#) or [Novirin](#).” - Greg Bennett, CBCD

In some cases, the antiviral drug acyclovir may cause poisoning in the brain that can lead to “visual hallucinations and dysarthria (difficult or unclear articulation of speech that is otherwise linguistically normal) and dyskinesia (abnormality or impairment of voluntary movement) of the arms and legs.” (1) It should be noted that these cases involved intravenous antiviral therapy. However, patients who have taken acyclovir orally have also experienced nausea, and vomiting, among many other side effects. For example, the American Academy of Family Physicians (AAFP) notes that “Effective oral antiviral medications are available for the treatment of genital herpes ... (and all can cause) ... Nausea, vomiting, headache, diarrhea, vertigo, myalgia, rash. (2) Drugs.com states that side effects, such as nausea, etc. associated with the use of acyclovir, the most widely recognized antiviral drug against herpes, may go away as an individual's body becomes used to it. (3)

However, in addition to side effects, it also “has poor bioavailability and a short half-life, which necessitates frequent dosing.” (2) In other words, the body does not absorb the drug well, and an individual must take doses often for the drug to be effective.



formula of these natural antiviral products was shown to reduce HSV symptoms in two separate post-marketing clinical studies that followed FDA guidelines. Additionally, it was designed to help the immune system target the latent form of the virus.

Click to learn more about [herpes symptoms](#).

The formula of Gene-Eden-VIR and Novirin was tested by Hanan Polansky and Edan Itzkovitz from the CBCD. The studies showed that the Gene-Eden-VIR and Novirin formula is effective against the family of herpes viruses, including HSV-1 and HSV-2. The clinical studies were published in the peer reviewed, medical journal Pharmacology & Pharmacy, the first, in a special edition on Advances in Antiviral Drugs. Study authors wrote that, "individuals infected with (HSV-1 or HSV-2)...reported a safe decrease in their symptoms following treatment with Gene-Eden-VIR." (4) The study authors also wrote that, "We observed a statistically significant decrease in the severity, duration, and frequency of symptoms." (4)

Both products can be ordered online on the Gene-Eden-VIR and Novirin websites, here:

<http://www.gene-eden-vir.com>

and

<http://www.novirin.com>

Gene-Eden-VIR and Novirin are natural antiviral dietary supplements. Their formula contains five natural ingredients: Selenium, Camellia Sinensis Extract, Quercetin, Cinnamomum Extract, and Licorice Extract. The first ingredient is a trace element, and the other four are plant extracts. Each ingredient and its dose was chosen through a scientific approach. Scientists at polyDNA, the company that invented and patented the formula, scanned thousands of scientific and medical papers published in various medical and scientific journals, and identified the safest and most effective natural ingredients against latent viruses. To date, Gene-Eden-VIR and Novirin are the only natural antiviral products on the market with published clinical studies that support their claims.

Note: Novirin shares the same formula as Gene-Eden-VIR. The difference between the two is that Novirin has higher quality and more expensive ingredients.

While nausea and vomiting may sound bad enough, each of the anti-herpes drugs may cause even more nasty side effects. Some of these include tremors, visual and auditory hallucinations, confusion and coma, are neuropsychiatric side effects. For example, all of these have been associated with the use of Valacyclovir (Valtrex). (5)

What treatments are available for herpes infections?

"Two types of antiviral treatments against HSV are available: topical and oral. The treatments include penciclovir, acyclovir, famciclovir, and valaciclovir. However, their effectiveness is limited. For instance, a meta-analysis of five placebo-controlled and two dose comparison studies evaluated the effect of aciclovir, famciclovir or valaciclovir on symptoms. The meta-analysis showed that oral antiviral therapy decreases the duration and the associated pain of an outbreak by merely one day." (4) There are also the natural supplements Novirin and Gene-Eden-VIR, which have a formula that was designed to help the immune system target the herpes virus when it is in a latent form.

The CBCD reminds the public that Gene-Eden-VIR and Novirin are not a cure. However, two clinical studies, published in peer reviewed medical journals showed that their formula decreases symptoms

in infected individuals. Specifically, the formula of Gene-Eden-VIR and Novirin was shown to safely and effectively reduce herpes symptoms. Additionally, it was designed to help the immune system target the latent virus.

"We recommend that individuals who are infected with a latent herpes virus take Gene-Eden-VIR or Novirin." – Greg Bennett, CBCD

Find us on Facebook here: <https://www.facebook.com/GeneEdenVIR>

Follow us Twitter on here: @HananPolansky

All orders of these products are completely confidential, and no information is shared or sold to any third party. Privacy is assured.

References:

(1) Neurology Advisor - "Case Report: Neurotoxicity With Acyclovir In Dialysis Patient" - November 30, 2015

<http://www.neurologyadvisor.com/general-neurology/case-report-neurotoxicity-acyclovir-dialysis-herpes-zoster/article/456507/>

(2) JOHN G. BEAUMAN, MAJ, MC, USA, Evans U.S. Army Hospital, Fort Carson, Colorado "Genital Herpes: A Review" Am Fam Physician. 2005 Oct 15;72(8):1527-1534.

(3) Drugs.com - Acyclovir - Acyclovir Side Effects

(4) Polansky, H. Itzkovitz, E. Gene-Eden-VIR Is Antiviral: Results of a Post Marketing Clinical Study. Published in September 2013.

<http://www.scirp.org/journal/PaperInformation.aspx?PaperID=36101>

(5) Chaudhari D1, Ginn D. "Valacyclovir-associated neurotoxicity in peritoneal dialysis patients." Am J Ther. 2014 Nov-Dec;21(6):e215-7.

Greg Bennett
CBCD
585-250-9999
email us here

This press release can be viewed online at: <http://www.einpresswire.com>

Disclaimer: If you have any questions regarding information in this press release please contact the company listed in the press release. Please do not contact EIN Presswire. We will be unable to assist you with your inquiry. EIN Presswire disclaims any content contained in these releases.

© 1995-2015 IPD Group, Inc. All Right Reserved.

