

Stress of Open-Heart Surgery Reduces Patients' Vitamin D Levels, But Supplementation Before and After Surgery Helps

New study shows that giving heart patients vitamin D3 supplements before and after open-heart surgery reduced their vitamin D deficiencies.

SALT LAKE CITY , UT, USA, March 15, 2018 /EINPresswire.com/ -- The stress of open-heart surgery significantly reduces patients' vitamin D levels, but aggressive supplementation with vitamin D3, just before and after surgery, can completely eliminate the observed drop in vitamin D, researchers have found.

Researchers at the Intermountain Medical Center Heart Institute in Salt Lake City conducted a study called ASSESS-D, in which they discovered two key findings related to vitamin D levels and open-heart surgery: first, that open-heart surgery patients generally have low levels of vitamin D to begin with, and secondly, that stress associated with the surgery further reduced their vitamin D levels by the time they were discharged from the hospital.

“

Now that we know that stress from surgery causes vitamin D levels to drop, we want to continue our research and see if supplementing vitamin D levels will help prevent heart problems in the future.”

*J. Brent Muhlestein, MD,
Intermountain Medical Center
Heart Institute*



The stress of open-heart surgery significantly reduces patients' vitamin D levels, but aggressive supplementation with vitamin D3, just before and after surgery, can completely eliminate the observed drop in vitamin D, researchers have found.

The Intermountain Medical Center Heart Institute study showed that giving heart patients vitamin D3 supplements before and after surgery reduced those deficiencies. Patients who were given the supplement had normal levels of vitamin D shortly after the surgery.

This connection is important since a deficiency of vitamin D is associated with an increased risk of heart attacks, strokes, congestive heart failure, and other cardiovascular conditions, including high blood pressure and diabetes, according to J. Brent Muhlestein, MD, lead author of the study, and cardiovascular researcher at the Intermountain Medical Center Heart Institute.

“We’re gathering more evidence that vitamin D deficiency is strongly associated with heart disease and death,” said Dr. Muhlestein. “Our study was a mechanistic approach to try and figure out what

exactly is happening with vitamin D, and perhaps give us some information as we press forward with a randomized trial to find out the real answer to the question of the impact of vitamin D on heart disease and related problems.”

Results from the study were presented this week at the American College of Cardiology Scientific Session in Orlando.

In the ASSESS-D study, Dr. Muhlestein and his team studied 150 randomized elective open-heart surgery patients. Half of the patients received three daily 50,000-unit doses of vitamin D3 starting before their procedures. The other half were given a placebo. Patients were monitored for vitamin D levels and major adverse cardiovascular events throughout their hospitalization, and then again six months after their surgery.

Among the patients who received a placebo, vitamin D levels dropped significantly after open-heart surgery. Among those who received vitamin D supplementation, their vitamin D levels actually rose into the normal range.

“Now that we know that the stress from surgery causes vitamin D levels to drop, we want to continue our research and see if supplementing vitamin D levels will help prevent heart problems in the future, given our understanding that low levels of vitamin D can cause an increased risk for heart problems,” said Dr. Muhlestein.

The researchers from the Intermountain Medical Center Heart Institute’s next step is to evaluate the effect of vitamin D supplementation on future cardiovascular risk among patients who present with a heart attack.

In an ongoing study, called the Target-D Trial, patients who come in with a heart attack are randomized to receive long-term vitamin D supplementation or not. They are then followed to see if the supplements reduce their risk of a second heart attack or other complications.

“We need hard evidence, and we hope the Target-D trial will give us real randomized controlled outcomes data to figure out if it’s good to take vitamin D supplements if you’re a heart patient who has low vitamin D levels,” said Dr. Muhlestein.

Vitamin D, commonly known as the sunshine vitamin, is a hormone produced by the skin. It helps regulate more than 200 genes throughout the body. Vitamin D keeps abnormal cells from multiplying in breast and colon tissues, helps regulate blood pressure in the kidney, and helps regulate blood sugar levels in the pancreas.

The human body produces vitamin D on its own, particularly when you’re out in the sun. Because people today are spending less time outdoors, and using more sunscreen when they are outdoors, the entire American population has general deficiency of the vitamin. Other factors can lead to lower vitamin D levels, including a person’s weight, skin, pigmentation, sex, age, and where they live.

Other members of the research team include: John R. Doty MD; Donald L. Lappe. MD; Kirk U. Knowlton; Heidi T. May, PhD; Tyler Barker, PhD; John F. Carlquist, PhD; Kristin Konery; Shannon Inglet, PharmD; Ben Chisum, BS; Viet Le, PA-C; Oxana Galenko; and Jeffrey L. Anderson, MD.

The Intermountain Medical Center Heart Institute, part of the Intermountain Healthcare system based in Salt Lake City, is one of the premier cardiovascular centers in the country.

###

Jess Gomez
Intermountain Medical Center
801-507-7455
email us here

This press release can be viewed online at: <http://www.einpresswire.com>

Disclaimer: If you have any questions regarding information in this press release please contact the company listed in the press release. Please do not contact EIN Presswire. We will be unable to assist you with your inquiry. EIN Presswire disclaims any content contained in these releases.

© 1995-2018 IPD Group, Inc. All Right Reserved.