

National Suicide Prevention Week -- How Local Health Departments Work to Reduce Rural Suicide Rates

Question and Answers with Local Health Department Leader Carol Moehrle RN, BSN

WASHINGTON, DC, UNITED STATES, September 11, 2019 / EINPresswire.com/ -- According to a recent study in the Journal of the American Medical Association, suicide rates are rising and are 25 percent higher in rural areas than in urban areas. The [National Association of County and City Health Officials](#) (NACCHO), the voice of the nation's nearly 3,000 local health departments, approaches suicide as a public health issue. Here is a discussion of rural suicide issues with rural health department leader Carole Moehrle of Idaho. For the past 25 years, Ms. Moehrle has served as the district director for a five-county public health district in North Central Idaho that serves a population of approximately 110,000. Ms. Moehrle has been active in NACCHO since 1992, and served as NACCHO president in 2010-2011. This year Ms. Moehrle received NACCHO's highest honor, the Maurice "Mo" Mullet Lifetime of Service Award.



Carol Moehrle

NACCHO

National Association of County & City Health Officials

The National Connection for Local Public Health

Q. Ms. Moehrle, why is the distinction being made between rural and urban suicide rates?

Suicide is one of the top 10 leading causes of death in the United States. The fact that suicide occurs more frequently in rural areas is not new. However, recent CDC data shows the disparity between rural and urban areas has slowly widened in recent years. The gap in suicide rates between rural and urban areas grew steadily from 1999 to 2015. And since 2007, the gap began widening more quickly. The 2016 national age-adjusted suicide rate reached the highest level seen in decades, and rural America is sustaining the largest increases. The states with the highest suicide rates per 100,000 population include: Montana, Alaska, Wyoming, New Mexico, Utah, Nevada, Idaho, and Oklahoma. In contrast, the states with the lowest suicide rates include: District of Columbia, New Jersey, New York, Massachusetts, Maryland, Connecticut, California, Illinois.

Q. What are the differences and emotional catalysts for rural suicide?

One of the insights into rural suicide is that the stresses in rural communities are different than in urban areas. Many economically struggling rural areas are experiencing the largest losses to suicide. It's not the economics itself, but the associated stress that leads to physical or mental effects on health. Isolation is a catalyst for rural suicide. Isolation is due to the decreases in population density in our rural areas, but also isolation due to lack of transportation, lack of social support, and lack of access to health care. The stigma around mental health is a real factor in rural America, as well. Much of rural America is also designated as health professional shortage areas, with limited, if any, access to mental health and preventive services.

Q. Why should local public health be involved in mental health and behavioral health issues?

Local health departments are tasked more than ever with filling the gaps in behavioral health services in our communities. Over half of U.S. counties do not have a single social worker, psychologist, or psychiatrist. Most rural counties do not have federally qualified health centers with providers. However, all counties have a public health department. In rural and frontier areas, local health departments are often one of the only providers, which means we are called on to provide community mental health support and prevention even if we do not have staff with behavioral health training.

Q. What role can public health play in suicide prevention?

Local public health can change the trajectory for suicide. Suicide is a complex behavior and requires a public health approach to prevention that identifies broad patterns of suicide and suicidal behavior throughout a population. Suicide prevention efforts can include: strategies to reduce risk factors like access to lethal means, promoting resiliency in our population that helps rid the stigma of mental health, universal screening and linking people to treatment services, and linking existing data systems to suicide prevention efforts, which would allow for better assessment of long term outcomes and enhance understanding of the broader impacts of suicide prevention interventions.

Q. What would you like Congress and the Administration to understand about rural suicide prevention?

Suicide is a public health issue and is preventable. Behavioral health concerns, including suicide, are an important issue for local public health. Unfortunately, most local health departments lack the resources necessary to invest in these issues, and temporary grants alone cannot sustain the infrastructure and workforce necessary to really turn the tide for these challenging issues. Greater, sustained investment in local health departments is necessary to bolster the available workforce to respond to mental health issues and implement suicide prevention strategies.

###

About NACCHO

The National Association of County and City Health Officials (NACCHO) represents the nation's nearly 3,000 local governmental health departments. These city, county, metropolitan, district, and tribal departments work every day to protect and promote health and well-being for all people in their communities. For more information about NACCHO, please visit www.naccho.org.

Theresa Spinner
National Association of County and City Health Officials
+1 2027835551
[email us here](#)

This press release can be viewed online at: <http://www.einpresswire.com>

Disclaimer: If you have any questions regarding information in this press release please contact the company listed in the press release. Please do not contact EIN Presswire. We will be unable to assist you with your inquiry. EIN Presswire disclaims any content contained in these releases. © 1995-2020 IPD Group, Inc. All Right Reserved.