

Reduced inpatient and severe respiratory disease visits during COVID-19 in Wuhan, China


USA, January 31, 2024 /EINPresswire.com/ -- This study investigated the trends in hospital visits for respiratory diseases in Wuhan, China, spanning the years 2018 to 2021. We found hospital visits for respiratory diseases decreased during the [COVID](#) lockdown. After the lockdown, there was a significant and lasting reduction in inpatient and severe visits for respiratory diseases. Compared to other respiratory diseases, acute lower respiratory infections experienced the largest reductions after the lockdown.

In Wuhan, China, a stringent lockdown was implemented to contain the spread of COVID-19, transitioning to a normalised prevention and control strategy. Considering the shared transmission route through the respiratory tract, mask-wearing, social distancing, and personal hygiene have been linked to reduced transmission of respiratory pathogens other than SARS-CoV-2. However, there have been less investigation into the broader spectrum of respiratory diseases particularly in less developed countries.

In a new study (<https://doi.org/10.1016/j.glt.2023.12.001>) published in the KeAi journal Global Transitions, a team of Chinese researchers examined the trends in hospital visits for acute and chronic respiratory diseases between 2018 and 2021 using the national health insurance data from Wuhan.

“Hospital visits for all the investigated diseases decreased during the lockdown period,” shares first author of the study Xuemin Zhu from Tsinghua University. “While visits for control conditions rebounded to pre-pandemic levels in the later normalised prevention and control period, visits for respiratory disease exhibit a more sustained impact from the pandemic.”

Notably, inpatient and severe condition visits for respiratory disease remained reduced even



Notes: Blue dots represent the observed values. Black lines represent fitted values in the interrupted time series analysis model. Blue shades denote the 95% confidence interval of the fitted values. The red lines are the average hospital visits between 2018 and 2020. Sandy lines indicate the counterfactual scenarios assuming no COVID-19 pandemic. (A)-(C) represent the trend for respiratory disease-related hospital visits by visit type. (D)-(O) represent the trend for respiratory disease-related hospital visits by visit type and discharge diagnosis. Severe condition is defined as visits requiring ventilation, intubation, or admission to an intensive care unit

Weekly hospital visits per million population in Wuhan between 2018 and 2021 across visit types and discharge diagnoses.

after the lockdown was lifted. Guanqiao Li, corresponding author of the study and also from Tsinghua University, emphasised that this research holds implications for future outbreak preparedness, healthcare resource allocation, and respiratory disease prevention and management.

“During the lockdown, community-level NPIs and government policies, particularly stay-at-home orders and travel restrictions have impeded patients’ access to healthcare services, primarily explaining the substantial decrease in hospital visits for all investigated diseases. Our findings underscore the importance of proactive preparedness by healthcare sectors to effectively manage outbreaks while ensuring the continued provision of essential medical services,” Li explains.

The team noted that the decrease in inpatient and severe condition visits for respiratory diseases upon the return to normal daily life is likely a result of the wide and consistent mask-wearing, social distancing, and personal hygiene improvement. While these reductions cannot be attributed to any single factor, the personal-level NPIs may have effectively reduced the incidence of severe respiratory diseases. Further, the researchers believe that healthcare systems should bolster their resilience to ensure rapid adaptability to diverse healthcare needs in the ongoing presence of the pandemics.

DOI

10.1016/j.glt.2023.12.001

Original Source URL

<https://doi.org/10.1016/j.glt.2023.12.001>

Funding information

This study was funded by the National Natural Science Foundation of China (72374119), Tsinghua University Initiative Scientific Research Program of Precision Medicine (2022ZLA001) and Sanming Project of Medicine in Shenzhen (NO. SZSM202111001)

Lucy Wang

BioDesign Research

[email us here](#)

This press release can be viewed online at: <https://www.einpresswire.com/article/685145326>

EIN Presswire's priority is source transparency. We do not allow opaque clients, and our editors try to be careful about weeding out false and misleading content. As a user, if you see something we have missed, please do bring it to our attention. Your help is welcome. EIN Presswire, Everyone's Internet News Presswire™, tries to define some of the boundaries that are reasonable in today's world. Please see our Editorial Guidelines for more information.

© 1995-2024 Newsmatics Inc. All Right Reserved.